

### Controlled pollination

Controlled pollination is critical to ensure the desired hybrid is produced by allowing the male parent to pollinate the female parent only. This is done to maintain the hybrid's purity or homogeneity, since the hybrid is the desired product, not the result of unwanted hybrid or pureline recombination. Various techniques are employed to ensure this, including:

1. Physical isolation of the hybrid seed production plots
2. Bagging the inflorescences or flowers
3. Emasculation or the use of male sterility
4. Hand-pollination

Careful monitoring is required to ensure the male parent is transferred to the female parent at the right time to prevent unwanted recombination or hybridization.

### Maintenance of genetic purity

It is also necessary to maintain genetic purity in order to retain the vigor and uniformity of the hybrids. Purity is maintained in the following ways:

1. Sufficient isolation distance between varieties
2. Removal of off-types
3. Clean harvesting, handling, and storage
4. Certification and quality control

Thus, high genetic purity ensures that the farmers get high-quality hybrid seeds that will produce the expected results.

### Hybrid Seed Production Systems

Different crops require different systems of hybridization due to their different requirements and goals of breeding. The system of hybridization is chosen according to the flower type, pollination pattern, economic viability, and availability of resources.

### Hand Emasculation and Pollination

This is the traditional method of hybridization and is commonly employed for crops that have bisexual flowers.

### Process:

1. The anthers are removed from the female parent before the emission of pollen grains
2. The emasculated flower is

### Male Sterility System

Male sterility is a natural or induced condition in which plants fail to produce functional pollen. This prevents self-pollination and eliminates the need for manual emasculation, making hybrid seed production more economical and efficient.

### Types of Male Sterility

#### a) Genetic Male Sterility (GMS)

The sterility is caused by nuclear genes. The sterility trait is inherited, and the sterility is expressed over generations in the offspring.

#### b) Cytoplasmic Male Sterility (CMS)

The sterility is caused by the action of mitochondrial genes, but it is maintained by the action of maintainer and restorer lines. CMS is commonly used in the production of hybrid seeds, as it is stable and easy to maintain.

#### c) Environment-Sensitive Male Sterility (ESMS)

The sterility is triggered by environmental factors, e.g., temperature or photoperiod. This is commonly used to develop two-line hybrid breeding programs.

Some crops commonly used for male sterility breeding programs are onion, cabbage, carrot, cauliflower, etc.

### Advantages:

1. No emasculation required
2. Cost reduction
3. Used for large-scale hybrid seed production

### Disadvantages:

1. Requires the maintenance of sterile lines
  2. Requires technical know-how
  3. Unstable conditions can cause problems with the sterility system
- Common crops: Onion, cabbage, carrot, and cauliflower.

### Self-Incompatibility System

Self-incompatibility is a genetically controlled mechanism that acts as a check against self-fertilization and encourages cross-pollination in plants. It is controlled by particular S-locus genes that control the interaction between pollen and stigma in plants. If the S-allele in the pollen is the same as that in the stigma, then the growth of the pollen tube is inhibited, preventing selfing.

### INTRODUCTION

The production of hybrid vegetable seeds is a specialized, skillful, and scientifically managed process aimed at producing high-quality crop varieties through the hybridization process. The main aim of hybridization is to take advantage of the heterosis phenomenon, whereby the first filial generation produced is better than the parent stock in terms of yield, uniformity, growth rate, adaptability, and resistance to diseases or environmental factors. This technology is considered one of the key technologies in the cultivation of vegetables in modern times, given its potential to develop crop varieties that can cope with the growing demand for high productivity, quality, and adaptability to environmental changes caused by climate change.

The production of hybrid seeds is carried out through scientifically managed breeding programs, where the main activities involve the production of parent lines, maintaining the purity of the parent lines, synchronizing the flowering times, controlling the pollination process, and providing isolation conditions. Unlike open-pollinated varieties, hybrid seeds require specialized technical know-how, specialized infrastructure, including isolation fields, pollination control methods, and quality testing protocols. For this reason, the production of hybrid seeds is carried out by experts rather than ordinary farmers.

The importance of hybrid vegetable seeds has been rising at a rapid rate over the last few decades due to their potential to improve the profitability of farming activities, along with the assured availability of fresh, processed, and export vegetables. Hybrids offer the advantage of uniformity in the size, color, and shape of the vegetables, making it easy to market the products. Moreover, some hybrid vegetables offer resistance to certain diseases, reducing the need to use chemical pesticides, thereby promoting the practice of integrated pest management in farming activities.



The promotion and advancement of hybrid vegetable technology across the globe have been highly supported by the leading research, development, and promotion organizations, including the Indian Council of Agricultural Research and the World Vegetable Center. These organizations are playing a vital role in the development of improved parental lines, promotion of hybrid technology, and strengthening the seed sector in developing, as well as developed, nations. The collaborative efforts of these organizations, along with national agricultural universities, private sector companies, and extension agencies, have contributed to the promotion of hybrid vegetables, thereby improving productivity, nutritional security, and income levels.

### Principles of Hybrid Seed Production

The foundation for hybrid seed production is built on basic genetic principles, which provide the guidelines for the production of high-quality F1 hybrids. These principles are the foundation for hybrid breeding programs, which define the quality, homogeneity, and yield potential of the produced hybrid seeds.

### Selection of the parental lines

The selection of the parental lines is the most critical factor in hybrid seed production. The selected parents should be:

1. Genetically pure or homozygous, to ensure the stable transmission of the desired characteristics
  2. Genetically diverse or contrasting, to ensure the highest heterosis
  3. Agronomically superior, to ensure the presence of the desired characteristics, including disease resistance, high yield potential, quality, and adaptability
- Several generations of selfing, selection, and evaluation are required to develop the parents to the desired level of homogeneity or purity before hybridization occurs.

# एग्रीकल्चर फ़ोरम फॉर टेक्निकल एजुकेशन ऑफ़ फार्मिंग सोसायटी

कोटा, राजस्थान



## Hybrid Vegetable Seed Production Techniques

संकलन

Shiv Narayan Dhaker<sup>1</sup>, Dr. Pramila<sup>2</sup>,  
Dr. Amit Kumar Singh<sup>3</sup>, Kalpana Kumar<sup>4</sup>

<sup>1</sup>Ph.D., Scholar, Division of Vegetable Science, ICAR- Indian Agricultural Research Institute, New Delhi (110012).

<sup>2</sup>Associate Professor, Department of Horticulture, Post-Graduate college of Agriculture, RPCAU, Pusa, Samastipur, Bihar (848125).

<sup>3</sup>Assistant Professor, Department of Post-Harvest Technology, College of Horticulture, Banda University of Agriculture and Technology, Banda (UP), (210001).

<sup>4</sup>Division of Fruit and Horticulture Technology, ICAR- Indian Agricultural Research Institute, New Delhi, (110012)

### There are two types of self-incompatibility:

**Gametophytic SI (GSi):** In this type, pollen compatibility is determined by the haploid genotype of the pollen itself.

**Sporophytic SI (SSi):** In this type, pollen compatibility is determined by the diploid genotype of the parent plant that produces the pollen.

Sporophytic SI is usually seen in Cole crops such as cabbage and cauliflower. This trait is useful in plant breeding because it can be used to produce hybrids without emasculation. Natural pollinators such as bees transfer pollen from one plant to another, ensuring cross-pollination between genetically different plants. This is important because self-incompatibility can sometimes be overcome in stressful conditions, due to mutation, and high temperatures.

#### 4. Gynoecious Line System

Gynoecious lines are lines that produce only female flowers, either due to genetic engineering or selection. Female flowers are the source of fruits and seeds, and these lines are useful in maximizing the production of fruit and seeds in hybrid varieties pollinated by the male parent. This method is more useful in cucurbits like cucumber and bitter gourd because:

- The flowering pattern of female flowers maximizes the efficiency of pollination.
- The cost of removing male flowers is eliminated.
- Uniformity of fruits in shape and size is improved.

To maintain gynoecious lines, male flowers must be induced for the purpose of producing seeds. This is done by applying silver nitrate or silver thiosulfate solutions, as these are ethylene inhibitors.

#### Development and Maintenance of Parental Lines

Hybrid seeds are produced by developing genetically pure parental lines through selfing and selection. Breeder seeds are the nucleus seeds maintained by plant breeders. These are kept under strict supervision. Foundation seeds are the seeds obtained by the multiplication of breeder seeds. These seeds are supplied to the producers of seeds.

#### Isolation Distance

Isolation prevents contamination from unwanted pollen sources. The distance requirement depends on pollination behavior and pollen dispersal ability. Wind-pollinated crops generally require greater isolation than insect-pollinated crops.

#### Flowering Synchronization

For hybridization to be effective, both male and female parents should flower at the same time. If their flowering is not synchronized, seed set is reduced and yield is compromised. Some advanced methods include:

- Adjusting sowing time to match duration of growth.
- Differential fertilizer application (nitrogen accelerates growth).
- Precise irrigation management.
- Foliar application of growth regulators such as gibberellic acids and growth retardants.
- Using temperature-controlled polyhouses to ensure precise timing.

#### Pollination Management

Hybrid seed production is all about effective pollination. Pollination methods vary from crop to crop:

- Insect-mediated pollination: Introduce bee colonies (*Apis mellifera* or *Apis cerana*) at recommended density.
- Hand pollination: Used for hybrids in crops such as tomato, brinjal, and chili peppers for research-level hybrids.
- Mechanical vibration: In greenhouse-grown tomatoes, electric vibrators are used to shake pollen from male flowers to ensure pollination.
- Male sterility: Cytoplasmic and genetic male sterility eliminate the need to emasculate flowers.

#### 5. Roguing

Roguing is practiced at different stages to eliminate off-types. Off-types could be genetic segregates, mechanical mixtures, and mutations. The different stages include:

- Seedling stage: Off-types are eliminated.
- Vegetative stage: Off-types are rogued out.
- Flowering stage: Off-types are rogued out.
- Fruiting stage: Off-types are rogued out.

Roguing ensures varietal purity, uniformity, and conformity to the norms of the certifying agency.

#### Harvesting and Seed Extraction

Hybrid fruits must be harvested at physiological maturity to ensure the development of well-formed embryos. Immature seeds will not germinate well.

#### Seed Extraction Methods:

- Wet Extraction: Used for tomato, cucumber, and melon. Fermentation is done to facilitate easy separation.
- Dry Extraction: Used for okra, beans, and peas. The pods are dried before the seeds are extracted.
- Mechanical Extraction: Used for large-scale seed production to improve efficiency.

After the seeds are extracted, they are washed and dried either in the shade or through dryers. The seeds are then graded based on size and density.

#### Seed Processing and Storage

Seed processing helps to maintain the viability of the seeds. The seeds are dried to a safe level to prevent the growth of fungi. The seeds are then treated with fungicides or other agents to prevent the growth of fungi. The seeds are then stored in airtight containers lined with laminated foil to maintain the viability of the seeds.

#### Advantages of Hybrid Vegetable Seeds

The advantages of hybrid seeds are:

- Heterosis or hybrid vigor
- Uniform crop stand, which is ideal for machine harvest
- Better resistance to multiple diseases
- Better tolerance to stress conditions
- Market demand, where the hybrid crop produces better-looking vegetables, hence more marketable

The yield advantage can be up to 20-50% or more than open-pollinated varieties under ideal conditions

#### Constraints in Hybrid Seed Production

- The constraints in hybrid seed production are:
- Higher cost of maintaining the parent stock
- Need for technical expertise
- Use of pollinators or the need for controlled pollination
- Environmental conditions, which affect the synchronization of flowering
- Stringent certification requirements
- Intellectual property rights on the hybrid

#### CONCLUSION

Hybrid vegetable seed production is a highly developed technology that combines aspects of genetics, plant physiology, agronomy, and biotechnology. It requires careful planning, precise execution, and quality control at every step from parental line development to final storage of the seed. With food needs rising and arable land falling worldwide, hybrid technology is a promising answer to boost productivity per unit area.

To advance hybrid technology in the future, it is necessary to blend traditional breeding expertise with cutting-edge molecular biology, digital technology, and climate-resilient approaches. Improving seed certification processes, enhancing farmer access to quality hybrid seed varieties, and investing in R&D infrastructure are also important to boost agricultural sustainability, food security, and farmer incomes.