Reg. No.: COOP/2023/KOTA/201080/26/210

**AGRICULTURE FORUM FOR TECHNICAL EDUCATION OF FARMING SOCIETY**

**Kota, Rajasthan**

# Your Soil Report, Your Fertilizer Plan: A Step-by-Step Guide

## Authors

**Rewangini Ranjha**

Department of Soil Science and Water Management
Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.

**Leaflet History**

Received : 13.10.2026
Revised : 18.10.2026
Accepted : 23.10.2026

## INTRODUCTION

A soil report really matters when the numbers tell you what to apply, how much of it to apply, and when to apply. A procedure of buying the same fertilizer you buy every other season for many of years can result in some nutrients becoming deficient while others may accumulate in the soil. In this way, soil testing shifts this type of practice to one plan based on the needs of your field, crop, and water availability. Still, a report is not a full diagnosis of all the problems that can limit production: poor drainage, compaction, pests, and unsuitable irrigation water drop growth. This could help guide you to documenting a written schedule of fertilizers from your sampling. The objective is balanced nutrition, dependable yields and better returns while protecting the productive capacity of your soil over the coming seasons.




### 2. Sample each Area

Divide the farm into sections with similar soil, slope, drainage and management history. Take samples from each section separately so strong and weak areas aren't combined. Walk in a zigzag across each section, collecting several small samples at the depth your lab recommends. Clear away loose plant material, but don't remove the soil beneath it. Skip unusual spots such as fertilizer piles, field edges, compost areas and irrigation channels or sample those separately if needed. Use clean equipment and a clean container. Mix the small samples well, then send the amount requested by the lab. A composite sample collected across the area gives a more reliable picture than soil taken from just one convenient spot.

### 3. Label Samples Properly

Collect samples before applying fertilizer or manure allowing time to get advice before sowing. Follow instructions by the laboratory on sampling depth, drying, packaging and delivery; special tests (mineral nitrogen may be one of them) may require prompt cooling or different handling. It is common demand that standard fertility samples are dried by air in shade, but this is not relevant to every analysis. Notate the samples with the name of the farmer, field identity, collection date, sample depth and chosen crop. Mention the previous crop, recent manure and fertilizer applications, irrigation source and any abnormalities of the field. The laboratory uses this information to interpret results appropriately. A field map (simple to this day) can be prepared such that similar areas can be defined for future samples and over time testing can be compared meaningfully.

### 4. Check the Units

Before you interpret the nutrient values, make sure that the investigation is a field study that you did yourself and that it covered the correct crop, sample date, and sampling depth. Note the units next to each result. Organic carbon may be shown as a percent, micronutrients as mg/kg, and available nutrients as kg/ha. The milligram per kilogram levels is

numerically equal to the parts per million system, but both measures do not measure how many kilograms of fertilizer to buy. Check if the available phosphorus and potassium are elements or oxide-related. Many fertilizer recommendations are expressed as N, $P_2O_5$ and $K_2O$. Never subtract a soil test number directly from crop nutrient demand: laboratory measurements require locally calibrated interpretation before they can guide a reliable fertilizer recommendation for planting.

**5. Check Soil Constraints**

Consider the reaction of the soil (as pH) with electrical conductivity, EC, before you determine the amount of fertilizer. Soil pH affects nutrient availability, and EC reflects the soluble salts based on the method used in the laboratory. Each of these should be read in the context of crop and test methodology. Soil acidity needs liming, but the quantity required depends on the measured liming requirement and the quality of the soil material. The application of gypsum is only justified if high pH is accompanied by increases in the sodicity that require quantification through appropriate testing of the solution. Salinity management may use drainage, appropriate irrigation water and regulated leaching where practicable. Increasing fertilizer applications on a salt affected field can cause even more stress; these constraints must be ameliorated with local technical guidance before raising nutrient applications.

**6. Interpret Nutrient Ratings**

Next, review the organic carbon result, nutrient ratings and any notes from the lab. Labels such as low, medium and high describe the likelihood of a crop responding to added nutrients within a particular



interpretation system. Their meaning can vary by laboratory, testing method and region, so don't treat them as universal standards. Low organic carbon points to a need for ongoing organic matter management, but it doesn't tell you how much manure to apply. A high phosphorus rating may support cutting back on phosphorus fertilizer, following local recommendations, even if another nutrient needs attention. Look at the report as a whole, not just the nitrogen result. Balanced nutrition means addressing real deficiencies not adding nutrients simply because they come in a familiar fertilizer blend.




**7. Match the Crop**

Adviser relevant information about your crop variety, planting period, irrigation availability and realistic potential yield. There are recommendations for your area, soil type and production system. They align fertilizer requirements with soil measures and achievable yield outcomes, based on results from many scientific investigations applying Soil test crop response (STCR) techniques. But field site data cannot be transferred as a rule of thumb from one site to another. Unattainable expectations can cause needless use of fertilizer when water is the limiting factor. The local Krishi Vigyan Kendra (KVK) or the agriculture university will help you convert the report



into nutrient rates. Make sure you consider if these rates account for organic inputs before making further assumptions during planning.

**8. Credit Existing Inputs**

List nutrient sources already-on hand to crop, including farmyard manure, compost, crop residues and any historically recognised prior legume credit for the site. Only a fraction of the total N in organic material becomes accessible in the first season, so overall nutrient content is not an immediate substitute for total fertilizer. Get a manure analysis when you can because nutrient contents of manure are dependent on diet, type of bedding, storage method and moisture content. Farmers should decide, in consultation and with direction from the adviser, how much is realistically per se available and subtract credits just once. There is normally also some supply of P and K in the manure, and double applications can lead to overbuild. Organic inputs improve soil condition but usage should build upon a tissue test nutrient budget, not be an automatic or default dose.

**9. Calculate Product Quantities**

Convert the recommended nutrient dose into the weight of fertilizer product using its guaranteed nutrient percentage. The basic calculation is fertilizer required equals nutrient required multiplied by 100, divided by nutrient percentage. For example, supplying 46 kilograms of nitrogen entirely through urea containing 46 percent nitrogen requires 100 kilograms of urea. This calculation illustrates conversion only; it is not a crop recommendation. Read the actual bag label because products differ in composition. Compound fertilizers supply more than

one nutrient, so count every contribution before selecting additional products. Compare suitable sources by the nutrients needed, application method and total field cost. The cheapest bag is not necessarily the most economical way to meet the recommendation.

**10. Work an Example**

Consider an illustrative recommendation of 120:60:40 kilograms N:$P_2O_5$:$K_2O$ per hectare, after all agreed nutrient credits. Using DAP containing 18 percent N and 46 percent $P_2O_5$, the phosphorus requirement needs approximately 130.4 kilograms DAP. This also supplies 23.5 kilograms nitrogen, leaving 96.5 kilograms nitrogen to provide. At 46 percent nitrogen, approximately 209.8 kilograms urea supplies that balance. Muriate of potash containing 60 percent $K_2O$ requires approximately 66.7 kilograms to supply the potassium. These quantities demonstrate arithmetic, not a prescription for any particular crop. Source suitability, chloride sensitivity and other nutrient needs still require checking. Most importantly, ignoring nitrogen supplied by DAP would overestimate the urea requirement and increase unnecessary nitrogen application to the field.

**11. Adjust for Area**

Convert the hectare recommendation to the actual cultivated area before purchasing fertilizer. Multiply kilograms per hectare by field area in hectares; one acre is approximately 0.4047 hectare. For a 500 square metre plot, the area is 0.05 hectare, so multiply each hectare quantity by 0.05. Using the illustrative calculation, this gives about 6.52 kilograms DAP, 10.49 kilograms urea and 3.34 kilograms muriate of potash for that plot over the planned season. These are product weights, not nutrient weights. Measure the planted area accurately and avoid relying on local land units unless their conversion is confirmed. Purchase according to the actual bag weight, then weigh application quantities instead of estimating them by handfuls or scoops.

**12. Schedule each Application**

Distribute the seasonal fertilizer total in ways that conform to crop demand and local recommendations. Split nitrogen applications are often advantageous in that a large initial dose may be lost before roots can use it. However, the correct location of phosphorus applications matters since it is relatively immobile in soil, while scheduling potassium treatments as well as the type of potassium fertilizer depends on crop type, texture of the soil and irrigation. Limit fertilizer uses near seed or young roots where it potentially could be harmful. Coordinate applications with adequate soil moisture, irrigation, and weather, times when it is Least likely to cause run off or major losses. Use compatible soluble products for fertigation and meet irrigation system requirements. For each planned application, include product, amount, growth stage and placement method in a written schedule of the entire crop.




**13. Address Specific Deficiencies**

Check sulphur and micronutrient results independently instead of just assuming that an NPK fertilizer covers all needs. Add zinc and boron and other micronutrients only if testing and locally proven crop guidance indicates use. Need may be small, and excessive application can ruin crops. Certifying the real material before computing its dose is essential, product concentration and chemical form being the drivers. Yellowing of the leaf alone can be misleading with regards to diagnosing the deficient nutrient because there is other could look alike symptoms such as waterlogging, root injury, disease and several deficiencies. Plant tissue analysis, properly sampled at the appropriate time, can be of help in finding causes for confusion. These foliar cuisines can be ductile deficiencies but will not replace the crop-wide program of soil fertility and nutrient management.

**14. Monitor Crop Response**

After application and to investigate patches before you splash more fertilizer, monitor crop growth. When suspecting nutrients, you should also consider checking for soil moisture problems, root stresses, insect pests, and drainage issues. When it is relevant, study the true nitrogen decision during the season by using validated leaf colour charts, crop sensors or tissue tests. They have to be crop specific, as either a numbers from soil test or using Digital imaging can help determine a dose of fertilizer. As research and extension programs increasingly combine soil testing with field observations and the right source, rate, time and place of application. This process will lead you to fine-tune your nutrient recommendations. Record any change and its reason, including additional product quantities, so the final nutrient total remains clear and avoidable overapplication is prevented.

**15. Record and Review**

Finish with a single field record that has the Soil report: including Crop, area, nutrient recommendation, inputs, products chosen and application dates Adding actual fertilizer purchases Labour cost irrigation events Yield Quality observations after harvest. These records verify whether it caused planned results or confounded ones. As often recommended in your local area repeat soil test all the same depths, seasons, methods; maybe do it to identify a specific condition or intensive cropping. Take the report and records to a trained adviser before the next purchase. The practical message is simple: sample accurately, interpret locally, calculate carefully, apply appropriately and review results. Your soil report becomes useful when it guides decisions throughout the farming season.

## CONCLUSION

Soil test data is a scientific foundation of balanced fertilizer management, better crop productivity, and sustainable soil health. Accurate nutrient interpretation, correct sampling, and crop-specific fertilizer recommendations can optimize nutrient use, reduce costs and lower environmental losses. Combining organic components, timely fertilizer applications, frequent monitoring and periodic soil testing to further improve nutrient use efficiency and soil fertility in the long term. Ultimately, proper agricultural practices which apply the correct dose of fertilizer at the proper time in the appropriate place are the key to efficient, productive and sustainable farming.